

TWO RAPIDLY VARIABLE GALACTIC X-RAY TRANSIENTS OBSERVED WITH *Chandra*, XMM AND *Suzaku*C. O. HEINKE<sup>1</sup>, J. A. TOMSICK<sup>2</sup>, F. YUSEF-ZADEH<sup>3</sup>, J. E. GRINDLAY<sup>4</sup>*Draft version June 25, 2009*

## ABSTRACT

We have identified two moderately bright, rapidly variable transients in new and archival X-ray data near the Galactic center. Both objects show strong, flaring variability on timescales of tens to thousands of seconds, evidence of  $N_H$  variability, and hard spectra. XMMU J174445.5-295044 is seen at 2-10 keV fluxes of  $3 \times 10^{-11}$  to  $< 10^{-12}$  ergs cm<sup>-2</sup> s<sup>-1</sup>, with  $N_H$  at or above  $5 \times 10^{22}$  cm<sup>-2</sup>, by XMM, *Chandra*, and *Suzaku*. A likely 2MASS counterpart with  $K_S = 10.2$  shows colors indicative of a late-type star. CXOU J174042.0-280724 is a likely counterpart to the fast hard transient IGR J17407-2808. *Chandra* observations find  $F_X(2-10 \text{ keV}) \sim 10^{-12}$  ergs cm<sup>-2</sup> s<sup>-1</sup>, with large  $N_H$  variations (from  $2 \times 10^{22}$  to  $> 2 \times 10^{23}$  cm<sup>-2</sup>). No 2MASS counterpart is visible, to  $K_S > 13$ . XMMU J174445.5-295044 seems likely to be a new symbiotic star or symbiotic X-ray binary, while CXOU J174042.0-280724 is more mysterious, likely an unusual low-mass X-ray binary.

*Subject headings:* binaries : X-rays — stars: neutron

## 1. INTRODUCTION

The wide field of view, sensitivity to hard X-rays, and program focused on the Galactic Plane of the INTEGRAL IBIS telescope has unveiled new kinds of hard X-ray transient behavior (Sguera et al. 2005). Many of these systems are high-mass X-ray binaries containing neutron stars (NSs), either eccentric Be systems where the NS feeds from an excretion disk, or supergiant fast X-ray transient systems (Negueruela et al. 2006) where the NS feeds from a (clumpy) stellar wind (Walter & Zurita-Heras 2007). A number of new INTEGRAL sources have now been identified with supergiant fast X-ray transients, showing rapid flaring behavior and variable  $N_H$ , and in some cases (slow) pulsations (Patel et al. 2004; Walter et al. 2006; Chaty et al. 2008).

However, some rapidly variable X-ray transient behavior may be produced by other systems, such as symbiotic stars (with white dwarf accretors, e.g. Smith et al. 2008), symbiotic X-ray binaries (with neutron star accretors, e.g. Masetti et al. 2007), some unusual black hole X-ray binaries (in't Zand et al. 2000), or magnetars (Castro-Tirado et al. 2008; Stefanescu et al. 2008). Identifying the optical/infrared counterparts of rapidly variable X-ray transients is important for understanding these systems, and requires accurate positions from focusing X-ray instruments.

The INTEGRAL hard X-ray source IGR J17407-2808 was first identified by Kretschmar et al. (2004) as a hard X-ray flaring source, positionally consistent with the faint ROSAT source 2RXP J174040.9-280852. Sguera et al. (2006) described the X-ray activity as three fast (few minute timescale) bright (up to 0.8 Crab) flares, and refined the position to a 1.7' radius. A 23 keV

bremsstrahlung model was an acceptable fit to the 20-60 keV spectrum. Sguera et al. (2006) noted that IGR J17407-2808's outburst was unusually strong and brief compared to known supergiant fast X-ray transients, and refrained from ruling out alternative interpretations.

We have identified two spectrally hard, rapidly variable, X-ray transients in archival *Chandra*, XMM-Newton, and *Suzaku* data near the Galactic Center. Our likely counterpart to the INTEGRAL IBIS fast transient IGR J17407-2808 detected in 2004, CXOU J174042.0-280724, showed strong flaring activity in early 2007. X-ray transient behavior has not been previously identified from the other, XMMU J174445.5-295044, for which we identify hard X-ray flaring in late 2006 and early 2007.

## 2. XMMU J174445.5-295044

## 2.1. Data

We observed the supernova remnant G359.1-0.5 on Sept. 24, 2006 (ObsId 0400340101) for 42 ksec with XMM's EPIC camera, using two MOS CCD detectors (Turner et al. 2001) with medium filters and one pn CCD detector (Strüder et al. 2001) with a thin filter. (The time resolution was 2.6 s for the MOS cameras and 73.4 ms for the pn camera.) All data were reduced using HEASOFT 6.5<sup>5</sup> and SAS version 8.0.0<sup>6</sup>. The data were moderately affected by soft proton flares. For spectral analysis, we chose a compromise between retaining photon statistics and excluding background, using limiting 0.2-10 keV rates of 5, 6, and 40 counts/s for MOS1, MOS2, and pn cameras (respectively). This left 25.0, 26.2 and 23.4 kiloseconds of time for the MOS1, MOS2, and pn cameras. We excluded event grades higher than 12 for MOS, and higher than 4 for the pn. The brightest point source was so much brighter than the background that all data could be used to produce lightcurves (see below).

We used *Chandra* archival ObsIDs 658, 2278, 2289 (all with the ACIS-I imaging CCD array), and 2714 (using the HRC-I high resolution camera). The data were re-

<sup>1</sup> University of Alberta, Dept. of Physics, Room #238 CEB, 11322-89 Avenue, Edmonton AB, T6G 2G7, Canada; cheinke@phys.ualberta.ca

<sup>2</sup> Space Sciences Laboratory, 7 Gauss Way, University of California, Berkeley, CA 94720-7450, USA

<sup>3</sup> Northwestern University, Dept. of Physics & Astronomy, 2145 Sheridan Rd., Evanston, IL 60208, USA

<sup>4</sup> Harvard University, Dept. of Astronomy, 60 Garden Street, Cambridge, MA 02138, USA

<sup>5</sup> <http://heasarc.gsfc.nasa.gov/docs/software/lheasoft>

<sup>6</sup> <http://xmm2.esac.esa.int/sas/>




FIG. 1.— Upper panel: Spectra of XMMU J174445.5-295044 from 2006, with XMM EPIC pn (top) and MOS (below; red online), fit with an absorbed powerlaw (Table 2). Lower panel: residuals to the fit.

duced with CIAO 4.1 and CALDB 4.1, following standard data analysis threads<sup>7</sup>. We used archival XIS data from Suzaku ObsID 501056010, reduced following the Suzaku Data Reduction Guide<sup>8</sup>. We used the cleaned event files in the HEASARC archive, extracted spectra and backgrounds with XSELECT<sup>9</sup>, and produced response files with the *xisresp* task. Finally, we included archival XMM data from three previous observations of 1E 1740.7-2942, in 2001, 2003, and 2005. These observations were conducted with the pn camera in timing mode, so only the MOS detectors were useful for our imaging. We performed standard filtering, as discussed above. All of these data are described in Table 1.

## 2.2. Discovery of XMMU J174445.5-295044

XMMU J174445.5-295044 is the brightest point source in the 2006 XMM data. We extracted its spectrum (from the data with high background excluded) and lightcurve (from the full dataset) from 20'' circles, with background regions nearby on the same chip.

XMMU J174445.5-295044 shows a very hard absorbed spectrum (Fig. 1, Table 2), indicating it is not a nearby coronal source. A power-law fit to the pn and combined MOS data is acceptable, and finds an intrinsically hard spectrum ( $\Gamma = 1.18^{+0.08}_{-0.07}$ ) with substantial extinction ( $N_H = 8.6 \pm 0.4 \times 10^{22} \text{ cm}^{-2}$ ). The spectrum is apparently featureless, with limits on the equivalent width of Fe lines at 6.4 and 6.7 keV at 26 and 9 eV respectively. Its average unabsorbed 2-10 keV X-ray luminosity would be  $1.7 \times 10^{35} \text{ ergs s}^{-1}$  if it were located at an 8 kpc distance.

We produced barycentered lightcurves for XMMU J174445.5-295044, and found strong flaring on timescales of hundreds of seconds (Fig. 2). The hardness of the spectrum does not vary strongly, indicating that the flaring is intrinsic and not due to varying  $N_H$ . Power spectra, with a binning timescale of 0.4 seconds, did not produce clear evidence of any periodicity, only red noise (rms variability of 24%, from  $2 \times 10^{-4}$  to 0.1 Hz).




FIG. 2.— 2006 XMM pn background-subtracted 1.5-9 keV lightcurve of XMMU J174445.5-295044, binned at 100s resolution. Below it (in red online), we plot the background (appropriately scaled) extracted from a nearby region on the same chip.




FIG. 3.— Images in 2MASS K band (left) and GLIMPSE 4.5 micron band (right) of the XMMU J174445.5-295044 field. Generous 4'' error circles are plotted around the XMM position, containing only one possible counterpart.

We use an error radius of  $2''^{10}$  for comparing the XMM position with accurate optical/IR catalogs (though we note that others, e.g. Walter et al. (2006), tend to use 4''). A potential 2MASS<sup>11</sup> counterpart (2MASS J17444541-2950446) is only 1.78'' from the XMM position, with  $K_S=10.2$  and  $J - K_S=4.7$  (Fig. 3, Tables 3, 4). Only 12 2MASS stars with magnitude  $K_S < 10.2$  lie within 50'' of XMMU J174445.5-295044, indicating that the probability of finding a star of this brightness within 2'' (our search radius) is only 2%. Thus, the 2MASS star is the likely counterpart of XMMU J174445.5-295044.

Since intrinsic  $J - K$  colors range from -0.2 for O stars to 1.2 for M giants,  $E(J-K)$  must be within 3.5 to 4.9. For standard interstellar reddening laws,  $A_V=20$  to 28, and  $N_H = 3.6 \times 10^{22}$  to  $5 \times 10^{22} \text{ cm}^{-2}$ . These values suggest that if this star is the true counterpart, the X-rays probably suffer additional local absorption.

We use the system of Negueruela & Schurch (2007) to compare the near-IR colors of the 2MASS counter-

<sup>7</sup> <http://cxc.harvard.edu/ciao/threads>

<sup>8</sup> <http://heasarc.gsfc.nasa.gov/docs/suzaku/analysis/abc/>

<sup>9</sup> <http://heasarc.gsfc.nasa.gov/docs/software/ftools>

<sup>10</sup> XMM-SOC-CAL-TN-0018, from <http://xmm.esac.esa.int>

<sup>11</sup> see <http://irsa.ipac.caltech.edu/>




FIG. 4.— 1.5-6 keV images of all 6 observations with detections or marginal detections of XMMU J174445.5-295044. All images are on the same scale (dimensions 6.3' by 8.7'), though the binning and stretch vary. Two observations (July 2001 and Sept. 2003) have been smoothed to more clearly identify XMMU J174445.5-295044. The Sept. 2003 MOS observation is only at best a marginal detection. Note that four of the observations place XMMU J174445.5-295044 near or at the edge of the detector.

part with those expected for high-mass X-ray binaries. Following Comerón & Pasquali (2005), we define the reddening-free parameter  $Q = (J - H) - 1.70(H - K_S)$ . Supergiants should have  $K_S < 11$  and  $Q < 0.2$ , Be stars should have  $K_S < 12$  and  $Q < 0$ , and typical late-type stars have  $Q \sim 0.5$ . We find  $Q = 0.81$  for the 2MASS counterpart, which is redder than expected even for late-type stars.

### 2.3. XMMU J174445.5-295044 at lower fluxes

The location of XMMU J174445.5-295044 ( $l=359.13$ ,  $b=-0.31$ ), at the edge of the *Chandra* wide Galactic Center survey field (Wang et al. 2002), and within 15' of the LMXB 1E 1740.7-2942, has been observed numerous times by X-ray satellites. We have searched the HEASARC archive<sup>12</sup> for detections or constraining upper limits. We report here archival detections of XMMU J174445.5-295044 at lower fluxes by Suzaku, XMM, and *Chandra*, and upper limits from *Chandra* and XMM (Table 2, Fig. 4). Observations by ROSAT, ASCA, EXOSAT, and Einstein do not detect it, but the upper limits are not particularly constraining, so we do not discuss those observations further.

XMMU J174445.5-295044 is consistent with the position of CXOGC J174445.5-295042 (Muno et al. 2006), seen in *Chandra* ObsID 2278. (The *Chandra* position is at the edge of a chip, so its positional uncertainty of 2.8", while consistent with XMMU J174445.5-295044, may not be fully trustworthy.) We have attempted spectral fitting to an absorbed power-law and absorbed bremsstrahlung




FIG. 5.— *Chandra* spectrum of XMMU J174445.5-295044, from 2000 data (ObsID 658), fitted with an absorbed powerlaw model (Table 2).

spectrum, using the unbinned data and the C statistic (Table 2). Testing the acceptability of a C-statistic fit is done through Monte Carlo simulations (we use 1000) of the model. The percentage of those simulations having a larger C-statistic than the data is reported as the 'goodness'; a high value (e.g. 95%) indicates a poor fit. Its 10-count detection at the very edge of the chip makes conclusions tentative.

A second *Chandra* ACIS-I observation (ObsID 658) identifies XMMU J174445.5-295044 on the off-axis S2 chip, with over 100 counts. We extracted a spectrum (binning the data with 10 counts/bin) and a lightcurve with 200s bins. Chi-square and KS tests found no evidence of variability in the lightcurve. We fit the spectrum (Fig. 5) with absorbed power-law and bremsstrahlung models (Table 2). Fitting the unbinned data with the C-statistic gave similar results (within the errors).

A third *Chandra* ACIS-I observation (ObsID 2289) did not detect XMMU J174445.5-295044, at the extreme corner of the off-axis S2 chip. Neither did a *Chandra* HRC-I observation where XMMU J174445.5-295044 was far off-axis (ObsID 2714). To determine upper limits for nondetections of XMMU J174445.5-295044 in this paper, we assume a spectrum similar to its faintest detections; an absorbed power-law with photon index 2 and  $N_H = 10^{23} \text{ cm}^{-2}$ . We extract the number of photons in a circle corresponding to the 50% encircled energy radius for photons of 4.5 keV at XMMU J174445.5-295044's off-axis angle, using an energy range of 1.5-6 keV. Using a larger nearby background region free of sources, we compute the expected number of background photons in this circle, scale it to the source region, and compute the 90% confidence lower limit (Gehrels 1986) on the number of expected background photons in the source region. The upper limit on the source count rate is derived by subtracting this lower limit from the observed photons, accounting for the encircled energy fraction and the vignetting of the telescope. Finally, we calculate flux upper limits using PIMMS<sup>13</sup>.

A recently archived *Suzaku* observation (ObsID 501056010, 2007) clearly detects XMMU J174445.5-295044 in all three operational XIS (Koyama et al. 2007) cameras (due to source confusion in the Galactic Center,

<sup>12</sup> <http://heasarc.gsfc.nasa.gov/db-perl/W3Browse/w3browse.pl>

<http://heasarc.gsfc.nasa.gov/db-perl/W3Browse/w3browse.pl>

<sup>13</sup> <http://asc.harvard.edu/toolkit/pimms.jsp>




FIG. 6.— *Suzaku* spectrum of XMMU J174445.5-295044, fitted with an absorbed powerlaw model (fit only above 2 keV). XIS1 data (diagonal crosses, red online) are lower above 4 keV than the combined XIS0 and XIS3 data (boxes, black online).




FIG. 7.— *Suzaku* combined XIS 1.5-6 keV lightcurve of XMMU J174445.5-295044, using 500s bins.

we do not consider the PIN data here). We extract spectra from a  $100''$  region of the cleaned data, combining the spectra from XIS CCDs 0 and 3, and binning the data at 40 photons/bin (Fig. 6). The variations in thermal emission from the Galactic Center and large PSF of *Suzaku* make background subtraction difficult below 2 keV. Since XMMU J174445.5-295044 shows little emission below 2 keV, we choose to ignore data below 2 keV for spectral fitting. The flux is a factor of 5 lower than observed by XMM, with a softer spectrum, and slightly higher  $N_H$ . We created a lightcurve from all detectors, in the energy range 1.5-6 keV, at 16 s binning. Variability, suggesting continued flaring, is best seen at 500 s binning (Fig. 7).

We retrieved three prior XMM observations of 1E1740.7-2942 from the archive (Table 1), taken in 2001, 2003, and 2005. In each case, the pn camera was in timing or small window mode, so we used only MOS data.




FIG. 8.— Mosaic of five *Chandra* observations in the 0.5-7 keV range, showing the large (green online) error circles for IGR J17407-2808 from Bird et al. (2007) and Sguera et al. (2006), and (red online) ellipses around detected sources. The candidate counterpart CXOU J174042.0-280724 is indicated by two black ticks. The small thick (green online) error circle indicates the location of 2RXP J174040.9-280852; note that no *Chandra* sources lie near it. SLX 1737-282 is the bright source producing a readout streak, near the bottom of the image.

The 2001 and 2003 data were heavily affected by background flaring, forcing us to use partly contaminated data. A combined MOS image of the 2001 data produced only an upper limit on the existence of XMMU J174445.5-295044. The 2005 data produced a clear detection of XMMU J174445.5-295044 (Fig. 4, lower left), with a flux comparable to the *Chandra* detections (Table 2). In the 2003 data, the position of XMMU J174445.5-295044 falls off the MOS2 chips, but lies on a MOS1 chip. Although the source is not detected with a blind detection algorithm, our upper limit method described above calculates  $11^{+2.2}_{-4.8}$  counts (at 90% confidence). Thus we consider this as a marginal detection (see Fig. 4, upper right).

### 3. CXOU J174042.0-280724: AN INTEGRAL SOURCE?

Sguera et al. (2006) give the revised INTEGRAL error circle as R.A.= $17^h40^m40^s.08$ , decl.= $-28^d08'24''$  (J2000), with error radius of  $1.7'$ . Bird et al. (2007) gives the error circle as  $17^h40^m42^s$ ,  $-28^d12.1'$ , with a  $4.2'$  error radius. The Galactic Bulge Latitude Survey (Grindlay et al., in prep.) covers both error circles thoroughly, between 4 overlapping 15 ks ACIS-I observations. We downloaded these four observations, plus the ACIS-S observation of Tomsick et al. (2008). We searched for periods of enhanced background, but found none. We created a mosaic image in the 0.5-7 keV energy band (Fig. 8).

The mosaic image shows eighteen sources in one or the other of the two overlapping error circles. (Note that no *Chandra* source is visible in the error circle for the ROSAT source 2RXP J174040.9-280852.) The brightest of these sources, CXOU J174042.0-280724 (indicated),




FIG. 9.— Spectrum of CXOU J174042.0-280724 in *Chandra* ObsID 8202, fit with an absorbed bremsstrahlung spectrum.

shows marked flaring variability and a hard spectrum showing strong variations in absorption, which suggest that this source is the soft X-ray counterpart to IGR J17407-2808. None of the other sources within the IGR J17407-2808 error circles show these behaviors, so we focus on CXOU J174042.0-280724. (Tomsick et al. (2008) identify the brightest source visible in Fig. 8 as the low-mass X-ray binary SLX 1737-282, which is not a likely counterpart for IGR J17407-2808.)

CXOU J174042.0-280724 is in the field of view of ObsIDs 8199, 8200 (on the edge of the chip), 8202, and 7526, all of which were observed between July 30 and August 1, 2007 (Table 1). The count rate is much higher in ObsID 8202 than the others, by almost a factor of 10. None of these datasets showed background flaring, so we used all the data, and extracted spectra and lightcurves using CIAO tools.

A power-law spectral fit of CXOU J174042.0-280724 in ObsID 8202 finds a hard, absorbed spectrum, with photon index  $\Gamma = 0.9^{+0.4}_{-0.6}$  and  $N_H = 1.7^{+0.7}_{-0.9} \times 10^{22} \text{ cm}^{-2}$  (Table 5; Fig. 9). A lightcurve (binned at 100 s) shows irregular sharp, short flares (Fig. 10a), reaching at least 100 times the lowest flux. No periodicities are detected in the overall power spectrum. Although the count rates between flares are low, there is some indication that part of the variability is due to variable absorption, as the emission is systematically softer during the brightest periods (Fig. 10).

We test this by extracting spectra from bright and faint parts of ObsID 8202, and fitting the (ungrouped) spectra with an absorbed power-law model, with photon index constrained between 0 and 3. Our spectral fit to the brightest 220 seconds of the first flare finds  $N_H = 0.8^{+1.5}_{-0.7} \times 10^{22} \text{ cm}^{-2}$ , and a 2-10 keV flux of  $F_X = 8^{+4}_{-2} \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$ . (We note that there is no evidence for pileup in the spectrum or image of this flare, due to the source's off-axis location). This is still three orders of magnitude fainter than the peak INTEGRAL flux from IGR J17407-2808. A similar fit to the data below 0.04 counts/s (excluding the last 5 ksec) gives  $N_H = 2.0^{+2.4}_{-1.6} \times 10^{22} \text{ cm}^{-2}$ , and a flux of  $F_X = 3.6^{+2.2}_{-1.3} \times 10^{-13} \text{ ergs cm}^{-2} \text{ s}^{-1}$ . This exercise




FIG. 10.— Top: Lightcurve of CXOU J174042.0-280724 in *Chandra* ObsID 8202, 0.5-7 keV, binned at 100 s resolution. Bottom: Hardness ratio of CXOU J174042.0-280724 in ObsID 8202, 4-8 keV counts over 1-4 keV counts, in 500 s bins.

strongly supports intrinsic source variation, and does not prove  $N_H$  variations during ObsID 8202.

CXOU J174042.0-280724's substantially smaller count rate in the other three observations is explained by comparison of the spectra from ObsIDs 8199 and 8202. The spectrum observed in 8199 is substantially harder than that in 8202, showing a low-energy cutoff near 4 keV vs. 1.5 keV (compare Fig. 9 vs. 11). Fitting bremsstrahlung models to both spectra confirms that the  $N_H$  value changed dramatically between the two observations, from  $N_H = 29^{+36}_{-7} \times 10^{22} \text{ cm}^{-2}$  to  $N_H = 2.2^{+0.5}_{-0.3} \times 10^{22} \text{ cm}^{-2}$ , while the unabsorbed 2-8 keV fluxes are consistent (Table 5). Thus the inter-observation variability may be ascribed entirely to varying  $N_H$ , although the flaring (present in ObsID 8199 also; Fig. 12) cannot. The remaining two observations (ObsIDs 8200 and 7526) offer less information (in 8200 the source lies on a chip edge, while 7526 was a shorter observation), producing spectra which are poorly constrained. However, the spectrum from ObsID 8200 is as hard as that from 8199 (only 3 of 27 photons are below 4 keV in ObsID 8200, compared to 7 of 63 in ObsID 8199), giving support to our explanation of the variability. The spectrum from ObsID 7526 is rather softer (7 of 19 counts below 4 keV), which might be due to intrinsic spectral evolution, or to  $N_H$  variations. Future observations could probe CXOU J174042.0-280724's spectral evolution.

The best *Chandra* position for CXOU J174042.0-280724 is RA=17:40:42.05±0.01, Dec=−28:07:24.6±0.1. Without astrometric matching of nearby X-ray sources to other wavelengths, we must add a systematic uncertainty of 0.6" (90% conf.). We have searched for possible counterparts in the 2MASS catalog, but the nearest star is 1.9" away, at a magnitude of 11.8. Based on the faintest 2MASS sources detected within 1', rough limits of  $J > 14$  and  $K_S > 13$  can be set for the error circle.




FIG. 11.— Spectrum of CXOU J174042.0-280724 in *Chandra* ObsID 8199, fit with an absorbed bremsstrahlung spectrum. Note the higher low-energy cutoff in comparison with Fig. 9.




FIG. 12.— Lightcurve of CXOU J174042.0-280724 in *Chandra* ObsID 8199, 0.5-7 keV, binned at 500s resolution.

No counterparts exist in SIMBAD or the USNO catalog.

#### 4. DISCUSSION

We show the fluxes, photon indices, and  $N_H$  measurements for XMMU J174445.5-295044 in Fig. 13. Clearly XMMU J174445.5-295044 spends most of its time at lower fluxes, below  $F_X = 3 \times 10^{-12}$  ergs cm<sup>2</sup> s<sup>-1</sup>. The 2006 XMM and 2007 *Suzaku* observations clearly show differences in photon index and  $N_H$ . The *Chandra* results also suggest a softening of the intrinsic spectrum, and increased  $N_H$ , at lower fluxes. The variable absorption and hard spectrum seem to rule out a coronal nature for the X-ray emission, suggesting an accreting compact object.

The evidence of rapid variability on timescales of minutes, the hard spectrum, and the intrinsic and variable absorption suggest a high-mass X-ray binary nature for XMMU J174445.5-295044 (White et al. 1995).




FIG. 13.— Fluxes, photon indices, and  $N_H$  measurements for XMMU J174445.5-295044 from constrained ( $0.5 < \Gamma < 3.0$ ) absorbed power-law spectral fits to XMM, *Chandra* and *Suzaku* data over 8 years (see Table 2).

However, the red colors of the infrared counterpart for XMMU J174445.5-295044 are inconsistent with a high-mass X-ray binary nature, suggesting a red giant companion. The spectrum and variability are similar to those of, e.g., 4U 1954+31 (Masetti et al. 2007), suggesting a symbiotic X-ray binary nature. However, some symbiotic stars also show hard spectra and rapid variability (Luna & Sokoloski 2007), so a white dwarf accretor is also a viable explanation. We note that there are strong similarities between accretion from the clumpy stellar wind of high-mass stars vs. that from giant stars onto a compact object. Indeed, several INTEGRAL sources with X-ray properties similar to high-mass X-ray binaries have recently been identified as symbiotic X-ray binaries (Masetti et al. 2007; Nespoli et al. 2008; Tomsick et al. 2008).

The substantial extinction to XMMU J174445.5-295044's infrared counterpart suggests a distance of at least a kpc, and thus X-ray luminosities of  $> 10^{32}$  ergs s<sup>-1</sup> in quiescence, and  $> 10^{33}$  ergs s<sup>-1</sup> at maximum. The flux range and strong variability of XMMU J174445.5-295044 suggest it may fit into the category of very faint X-ray transients (Wijnands et al. 2006; Degenaar & Wijnands 2008), although it may reach higher fluxes at other times. Inspection of the ASM and RXTE bulge scan data finds no evidence of outbursts from this source, with the bulge scan providing typical upper limits of 2-10 mCrab (C. Markwardt, priv. comm, 2008; R. Remillard, priv. comm, 2008). Some of the other known very faint X-ray transients may be symbiotic systems; this possibility might be ruled out with deep infrared observations.

CXOU J174042.0-280724 has a more mysterious nature than XMMU J174445.5-295044, due to its lack of an optical counterpart. Even if it were not associated with IGR J17407-2808 (which we think it is), its varying  $N_H$  and rapid variability suggest accretion onto a compact

object. The high flux attained by IGR J17407-2808 indicates a neutron star or black hole nature. The distance to this source is unknown, as we do not have a measurement of the interstellar reddening, and at least some of the observed (variable) absorption must be intrinsic. This direction ( $l=0.12$ ,  $b=1.35$ ) is sufficiently reddened ( $N_H$  up to  $1.4 \times 10^{22} \text{ cm}^{-2}$ , Marshall et al. (2006)) that interstellar reddening could explain the observed  $N_H$  in ObsID 8202, which has the lowest observed  $N_H$  value. The  $K_S$  limit implies  $M_K > -2.2$  for a distance of 10 kpc, excluding main sequence stars with spectral types earlier than B1 and giant stars with spectral types later than K2. At a distance of 15 kpc, these limits shift to excluding main sequence stars earlier than O9 and giants later than K4, while supergiants are ruled out for either distance.

Some low- or intermediate-mass X-ray binaries, such as V4641 Sgr (in't Zand et al. 2000; Revnivtsev et al. 2002; Chaty et al. 2003) or XTE J1901+014 (Smith et al. 2007; Karasev et al. 2007), show rapid flaring similar to that seen from IGR J17407-2808. Alternatively, this object could be a Be-type binary at a distance over 10 kpc. We note that for a distance of 10 kpc, the peak INTEGRAL flux corresponds to  $L_X(20\text{-}60 \text{ keV}) = 1.1 \times 10^{38} \text{ ergs s}^{-1}$ , suggesting a bolometric flux above the Eddington limit for a neutron star.

Further studies of these mysterious sources are strongly encouraged to improve our understanding of their nature and of fast X-ray transient behavior. The most critical need is for deep optical and infrared imaging to search for CXOU J174042.0-280724's expected donor star. Spectroscopy of such an object (if possible) would be very valuable. Optical and/or infrared spectroscopy of XMMU J174445.5-295044's optical counterpart is also needed, to identify its spectral type and surface gravity

and search for evidence of accretion. This will test the hypothesis presented here that XMMU J174445.5-295044 is a symbiotic system. If the distance can be constrained, the nature of the accreting object (white dwarf or neutron star?) may be determined. Additional X-ray observations of both objects will also be useful to monitor their long-term activity, especially if CXOU J174042.0-280724 produces additional outbursts detectable by INTEGRAL.

COH thanks Craig Markwardt and Ron Remillard for their rapid efforts to search the RXTE bulge scans and RXTE/ASM records, respectively. COH acknowledges support by NSERC and NASA XMM GOF funding. JAT acknowledges partial support from *Chandra* award number GO8-9055X issued by the *Chandra X-Ray Observatory Center*, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of the National Aeronautics and Space Administration (NASA), under contract NAS8-03060. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and IPAC/Caltech, funded by NASA and the NSF. This research has made use of the NASA/IPAC Infrared Science Archive, and the Spitzer Space Telescope, which are operated by the Jet Propulsion Laboratory, Caltech, under contract with NASA. This research has made use of data obtained from the High Energy Astrophysics Science Archive Research Center (HEASARC), provided by NASA's Goddard Space Flight Center. It is based in part on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA member states and NASA.

## REFERENCES

- Bird, A. J. et al. 2007, *ApJ Supp*, 170, 175  
 Bodaghee, A., et al. 2007, *A&A*, 467, 585  
 Castro-Tirado, A. J. et al. 2008, *Nature*, 455, 506  
 Chaty, S., Charles, P. A., Martí, J., Mirabel, I. F., Rodríguez, L. F., & Shahbaz, T. 2003, *MNRAS*, 343, 169  
 Chaty, S., Rahoui, F., Foellmi, C., Tomsick, J. A., Rodríguez, J., & Walter, R. 2008, *A&A*, 484, 783  
 Comerón, F., & Pasquali, A. 2005, *A&A*, 430, 541  
 Degenaar, N., & Wijnands, R. 2009, *A&A*, 495, 547  
 Gehrels, N. 1986, *ApJ*, 303, 336  
 in't Zand, J. J. M. et al. 2000, *A&A*, 357, 520  
 Karasev, D. I., Lutovinov, A. A., & Grebenev, S. A. 2007, *Astronomy Letters*, 33, 159  
 Koyama, K. et al. 2007, *PASJ*, 59, 23  
 Kretschmar, P., Mereghetti, S., Hermsen, W., Ubertini, P., Winkler, C., Brandt, S., & Diehl, R. 2004, *The Astronomer's Telegram*, 345, 1  
 Luna, G. J. M. & Sokoloski, J. L. 2007, *ApJ*, 671, 741  
 Marshall, J. D., Robin, A. C., Reylé, C., Schultheis, M. & Picaud, S. 2006, *A&A*, 453, 635  
 Masetti, N., et al. 2007, *A&A*, 464, 277  
 Munro, M. P., Bauer, F. E., Bandyopadhyay, R. M., & Wang, Q. D. 2006, *ApJ Supp*, 165, 173  
 Negueruela, I., Smith, D. M., Reig, P., Chaty, S., & Torrejón, J. M. 2006, *ESASP "The X-ray Universe 2005"*, ed. Wilson, 604, 165  
 Negueruela, I., & Schurch, M. P. E. 2007, *A&A*, 461, 631  
 Nespoli, E., Fabregat, J., & Mennickent, R. E. 2008, *The Astronomer's Telegram*, 1450, 1  
 Patel, S. K., Kouveliotou, C., Tennant, A., Woods, P. M., King, A., Finger, M. H., Ubertini, P., Winkler, C., Courvoisier, T. J.-L., van der Klis, M., Wachter, S., Gaensler, B. M., & Phillips, C. J. 2004, *ApJ*, 602, L45  
 Revnivtsev, M., Gilfanov, M., Churazov, E., & Sunyaev, R. 2002, *A&A*, 391, 1013  
 Sguera, V. et al. 2005, *A&A*, 444, 221  
 —. 2006, *ApJ*, 646, 452  
 Smith, R. K., Mushotzky, R., Mukai, K., Kallman, T., Markwardt, C. B. & Tueller, J., 2008, *PASJ*, 60, 43  
 Smith, D. M., Rampy, R. A., Negueruela, I., & Torrejón, J. M. 2007, *The Astronomer's Telegram*, 1268, 1  
 Stefanescu, A., Kanbach, G., Slowikowska, A., Greiner, J., McBreen, S., & Sala, G. 2008, *Nature*, 455, 503  
 Strüder, L. et al. 2001, *A&A*, 365, L18  
 Tomsick, J. A., Chaty, S., Rodríguez, J., Walter, R., & Kaaret, P. 2008, *ApJ*, 685, 1143  
 Turner, M. J. L. et al. 2001, *A&A*, 365, L27  
 Walter, R., et al. 2006, *A&A*, 453, 133  
 Walter, R. & Zurita-Heras, J., 2007, *A&A*, 476, 335  
 Wang, Q. D., Gotthelf, E. V., & Lang, C. C. 2002, *Nature*, 415, 148  
 White, N. E., Nagase, F., & Parmar, A. N. 1995, in *X-ray binaries*, p. 1 - 57, 1-57  
 Wijnands, R. et al. 2006, *A&A*, 449, 1117

TABLE 1  
X-ray observations of XMMU J174445.5-295044 and CXOU J174042.0-280724

| Mission/Instrument     | ObsID      | Date              | Exposure     | Notes              |
|------------------------|------------|-------------------|--------------|--------------------|
| XMMU J174445.5-295044  |            |                   |              |                    |
| <i>Chandra</i> /ACIS-I | 658        | 2000-08-30        | 9235         |                    |
| XMM/MOS                | 0112971701 | 2001-03-31        | 7003         | Nondetection       |
| <i>Chandra</i> /ACIS-I | 2278       | 2001-07-20        | 11611        | At chip edge       |
| <i>Chandra</i> /ACIS-I | 2289       | 2001-07-21        | 11611        | Nondetection       |
| <i>Chandra</i> /HRC-I  | 2714       | 2002-03-02        | 29709        | Nondetection       |
| XMM/MOS                | 0144630101 | 2003-09-11        | 4477         | Marginal detection |
| XMM/MOS                | 0303210201 | 2005-10-02        | 21823        |                    |
| XMM/MOS                | 0400340101 | 2006-09-24        | 41028(25026) | Source flaring     |
| XMM/pn                 | 0400340101 | 2006-09-24        | 35337(23378) | Source flaring     |
| Suzaku/XIS             | 501056010  | 2007-03-13        | 26545        | Variable           |
| CXOU J174042.0-280724  |            |                   |              |                    |
| <i>Chandra</i> /ACIS-I | 8199       | 2007-07-30, 01:31 | 14682        | Source flaring     |
| <i>Chandra</i> /ACIS-I | 8200       | 2007-07-30, 05:57 | 14566        | At chip edge       |
| <i>Chandra</i> /ACIS-I | 8202       | 2007-07-30, 14:41 | 14300        | Source flaring     |
| <i>Chandra</i> /ACIS-S | 7526       | 2007-08-01, 13:04 | 5108         |                    |

NOTE. — For ObsID 0400340101, we give the exposure with background flares removed in parentheses (these data are used for spectral analysis).

TABLE 2  
Spectral fits to XMMU J174445.5-295044

| Mission/Instrument     | ObsID      | Flux<br>ergs cm <sup>-2</sup> s <sup>-1</sup> | $\Gamma$               | $N_H$<br>10 <sup>22</sup> cm <sup>-2</sup> | $\chi^2/\text{dof}$  | kT<br>keV           | $N_H$<br>10 <sup>22</sup> cm <sup>-2</sup> | $\chi^2/\text{dof}$ |
|------------------------|------------|-----------------------------------------------|------------------------|--------------------------------------------|----------------------|---------------------|--------------------------------------------|---------------------|
| Power-law fit          |            |                                               |                        |                                            | -Bremsstrahlung fit- |                     |                                            |                     |
| <i>Chandra</i> /ACIS-I | 658        | $1.6^{+0.6}_{-0.7} \times 10^{-12}$           | $3.0^{+0.2}_{-1.1}$    | $15^{+3}_{-6}$                             | 0.68/10              | $1.8^{+5.0}_{-1.0}$ | $19^{+14}_{-9}$                            | 0.57/10             |
| XMM/MOS                | 0112971701 | $< 3 \times 10^{-13}$                         |                        |                                            |                      |                     |                                            |                     |
| <i>Chandra</i> /ACIS-I | 2278       | $2.8^{+1.5}_{-1.1} \times 10^{-13}$           | $0.5^{+2.5}_{-0.2}$    | $5.1^{+21}_{-2.7}$                         | 146/15 <sup>b</sup>  | $> 2.3$             | $11^{+8}_{-7}$                             | 146/85 <sup>b</sup> |
| <i>Chandra</i> /ACIS-I | 2289       | $< 5 \times 10^{-13}$                         |                        |                                            |                      |                     |                                            |                     |
| <i>Chandra</i> /HRC-I  | 2714       | $< 1.6 \times 10^{-12}$                       |                        |                                            |                      |                     |                                            |                     |
| XMM/MOS                | 0144630101 | $1.3^{+0.3}_{-0.5} \times 10^{-12}$           |                        |                                            |                      |                     |                                            |                     |
| XMM/MOS                | 0303210201 | $6^{+6}_{-2} \times 10^{-13}$                 | $1.8^{+1.2}_{-1.3}$    | $8^{+9}_{-5}$                              | 0.54/11              | $> 1.8$             | $7.8^{+9.5}_{-3.3}$                        | 0.54/11             |
| XMM/EPIC               | 0400340101 | $2.29 \pm 0.05 \times 10^{-11}$               | $1.18^{+0.08}_{-0.07}$ | $8.6^{+0.4}_{-0.4}$                        | 0.91/368             | $> 126$             | $8.8^{+0.2}_{-0.2}$                        | 0.92/368            |
| Suzaku/XIS             | 501056010  | $5.2^{+2.8}_{-1.2} \times 10^{-12}$           | $2.4^{+0.6}_{-0.6}$    | $16^{+5}_{-4}$                             | 0.65/111             | $6.6^{+11}_{-2.9}$  | $13.9^{+3.9}_{-3.2}$                       | 0.66/111            |

NOTE. — Photon indices are restricted to the physically reasonable range of 0.5 to 3. Fluxes are unabsorbed, for 2-10 keV, using best-fit absorbed power-law model, or a power-law spectrum with  $\Gamma = 2$  and  $N_H = 1 \times 10^{23}$  cm<sup>-2</sup> for upper limits.

<sup>a</sup>: Error calculation reached hard limit.

<sup>b</sup>: Used C-statistic, reporting C-statistic and “goodness” (see text) instead of  $\chi^2$  and degrees of freedom.

TABLE 3  
Positions of XMMU J174445.5-295044, CXOU J174042.0-280724, and possible counterparts

| Name                    | RA          | Dec         | $\Delta$     | Notes    |
|-------------------------|-------------|-------------|--------------|----------|
| XMMU J174445.5-295044   | 17:44:45.55 | -29:50:44.3 | $\pm 2.0^a$  | XMM pos. |
| XMMU J174445.5-295044   | 17:44:45.54 | -29:50:42.1 | $\pm 2.8^b$  | Muno+06  |
| 2MASS J17444541-2950446 | 17:44:45.41 | 29:50:44.6  | $\pm 0.07^a$ |          |
| CXOU J174042.0-280724   | 17:40:42.05 | -28:07:24.6 | $\pm 0.6^b$  | CXO pos. |

NOTE. — Positions of detected (X-ray or optical/IR) source and error. <sup>a</sup>: 1 $\sigma$  error.  
<sup>b</sup>: 90% confidence error.

TABLE 4  
Magnitudes and Fluxes of 2MASS J17444541-2950446

| Band                 | Magnitude | Flux  |
|----------------------|-----------|-------|
| <i>J</i>             | 14.89(5)  | 1.77  |
| <i>H</i>             | 11.62(4)  | 23.1  |
| <i>K<sub>S</sub></i> | 10.17(3)  | 57.2  |
| 3.6 $\mu$ m          | 9.06(5)   | 66(3) |
| 4.5 $\mu$ m          | 8.87(5)   | 51(2) |
| 5.8 $\mu$ m          | 8.67(4)   | 40(2) |
| 8.0 $\mu$ m          | 8.67(5)   | 22(1) |

NOTE. — Fluxes in mJy. Magnitudes from 2MASS and GLIMPSE.

TABLE 5  
X-ray Spectra of CXOU J174042.0-280724

| Mission/Instrument     | ObsID | Flux<br>ergs cm <sup>-2</sup> s <sup>-1</sup> | $\Gamma$             | $N_H$<br>10 <sup>22</sup> cm <sup>-2</sup> | $\chi^2/\text{dof}$ | $kT$<br>keV | $N_H$<br>10 <sup>22</sup> cm <sup>-2</sup> | $\chi^2/\text{dof}$ |
|------------------------|-------|-----------------------------------------------|----------------------|--------------------------------------------|---------------------|-------------|--------------------------------------------|---------------------|
| Power-law fit          |       |                                               |                      |                                            | Bremsstrahlung fit  |             |                                            |                     |
| <i>Chandra</i> /ACIS-I | 8199  | $7^{+24}_{-3} \times 10^{-13}$                | $0.5^{+2.5}_{-0a}$   | $24^{+37}_{-7}$                            | 1.5/3               | > 2.4       | $29^{+36}_{-7}$                            | 1.7/3               |
| <i>Chandra</i> /ACIS-I | 8200  | $2.1^{+2.2}_{-1.6} \times 10^{-12}$           | $3^{+0a}_{-2.5a}$    | $48^{+24}_{-27}$                           | 147/30 <sup>b</sup> | > 0.5       | $58^{+69}_{-20}$                           | 146/99 <sup>b</sup> |
| <i>Chandra</i> /ACIS-I | 8202  | $1.3^{+0.2}_{-0.2} \times 10^{-12}$           | $0.8^{+0.5}_{-0.3a}$ | $1.5^{+0.8}_{-0.6}$                        | 1.23/16             | > 25        | $2.2^{+0.5}_{-0.3}$                        | 1.39/16             |
| <i>Chandra</i> /ACIS-S | 7526  | $1.7^{+1.0}_{-0.8} \times 10^{-13}$           | $0.5^{+0.6}_{-0a}$   | $2.8^{+4.7}_{-1.7}$                        | 148/62 <sup>b</sup> | > 10        | $3.6^{+6}_{-1.8}$                          | 152/85 <sup>b</sup> |

NOTE. — Fluxes are unabsorbed, 2-10 keV, using best-fit absorbed powerlaw model (constraining photon index to range 0.5 to 3).

<sup>a</sup>: Error calculation reached hard limit.

<sup>b</sup>: Used C-statistic, reporting C-statistic value and “goodness” (see text) instead of  $\chi^2$  and degrees of freedom.